Detecting Active Fires using Himawari-8: a report from the NSW trial

Research advisory forum / 2019

Dr Chermelle Engel*/ RMIT University, Bushfire and Natural Hazards CRC Prof Simon Jones / RMIT University Assoc Prof Karin Reinke/ RMIT University Dr Stuart Matthews / NSW RFS Dr Alex Holmes* / NSW RFS

*Presenters





Business Cooperative Research Centres Programme





https://worldview.earthdata.nasa.gov. Aqua/MODIS (true color + fire anomalies)



Polar-Orbiting



Source: NOAA Science on a Sphere website

https://sos.noaa.gov/

Himawari-8



Source: Japan Meteorological Agency website http://www.jma-net.go.jp





IBRA

Interim Biogeographic Regionalisation for <u>Australia</u>

419 sub-regions





Work out likely MIR hotspots not associated with cloud





2019 mid-march to mid-July with break in middle







True Color

MIR (3.9 μm) 10:00 10/08/2019 AEST











"An operational user's view of hotspot markers overlaid on the most recent Himawari-8 #7 band image. Controls on the left allow temporal filtering of hotspots. Users can click on hotspot markers to display information about the detection."















- Linescan image 10:00 next morning



Polar & H8IBRA inter-comparison Within 1-pixel, within 10-minutes ...



Polar platform	Confidence, (FRP > 30 MW) (FRP > 0 MW)	%Polar found	%H8IBRA found
VIIRS	conf = (h or n)	81% (28%)	79% (94%)
Aqua	conf > 49	75% (36%)	83% (90%)
Terra	conf > 49	77% (26%)	89% (92%)

2019 mid-march to mid-July with break in middle



14/05/2019 @ 14:08 AEST



Wrap Up

- The trial helped to focus the research
- Invaluable tool for communication between stakeholders

Next steps:

- Further inter-comparison
- Expand trial into other seasons and regions within Australia?
- Aid rural fire services to adapt to such high-frequency data feed





QUESTIONS?

True Color



